

The Impact of Lavender Essential Oil on the Human Nervous System: A Review

Camila Castellan Cardoso ^{1,*}, Francisco Rafael Oliveira da Silva ¹

¹ Centro Universitário de Tecnologia de Curitiba, Curitiba, Paraná, Brazil.

* Correspondence: camiscas1@hotmail.com.

Abstract: Lavender (*Lavandula angustifolia*) has been traditionally used for its therapeutic properties, particularly in treating neurological and psychological disorders. Recent human studies suggest that lavender oil exhibits anxiolytic, mood-stabilizing, sedative, analgesic, and neuroprotective effects. This review examines the clinical evidence supporting lavender's efficacy in neurological applications, emphasizing its potential mechanisms and therapeutic benefits.

Keywords: Aromatherapy; Essential Oils; Human Nervous System; Lavender Essential Oil.

1. Introduction

Lavender, a plant native to the Mediterranean region, has been widely used for centuries for its medicinal and aromatic properties. Traditionally, it has been employed in herbal remedies, perfumes, and therapeutic applications. Historical records indicate that ancient civilizations, including the Romans, Greeks, and Egyptians, utilized lavender for its calming and healing properties, particularly in treating ailments related to the nervous system [1].

The primary constituents of lavender essential oil include linalool and linalyl acetate, which are believed to contribute significantly to its neurological effects. These compounds exhibit anxiolytic, sedative, and mood-stabilizing properties, which make lavender a popular natural alternative for managing stress-related conditions. Additionally, these bioactive components have demonstrated antioxidant and anti-inflammatory properties, which may further support neuroprotection and cognitive function [2, 3].

Over the past few decades, scientific research has increasingly focused on validating the therapeutic potential of lavender essential oil. Clinical studies have explored its efficacy in treating anxiety, depression, sleep disturbances, pain, and cognitive dysfunction. Various administration methods, including aromatherapy, oral consumption, and topical application, have been investigated to determine the optimal delivery route for its therapeutic benefits [4].

Aromatherapy, which involves inhaling lavender essential oil, has been shown to activate brain regions associated with relaxation and emotional regulation, such as the amygdala and hippocampus. This mechanism suggests that lavender may exert its effects through modulation of the limbic system. Studies utilizing functional neuroimaging have provided evidence that lavender influences neural activity patterns in a manner consistent with anxiolytic and sedative agents. Moreover, the widespread use of lavender in complementary and alternative medicine has spurred interest in its integration into conventional medical treatments [5, 6]. Some researchers have proposed that lavender could serve as an adjunct therapy for neurological and psychiatric conditions, potentially reducing the reliance on pharmaceutical interventions that often come with adverse effects.

Citation: Cardoso CC, Silva FRO. The Impact of Lavender Essential Oil on the Human Nervous System: A Review. Brazilian Journal of Aromatherapy and Essential Oil. 2025;2:bjhae22.

doi: <https://doi.org/10.62435/2965-7253.bjhae.2024.bjhae22>

Received: 1 September 2025

Accepted: 10 September 2025

Published: 16 September 2025



Copyright: This work is licensed under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

However, despite promising findings, further large-scale clinical trials are necessary to establish standardized dosing guidelines and long-term safety profiles [7-9].

Lavender's historical significance and modern scientific validation highlight its potential as a natural therapeutic agent for neurological health. As research continues to unravel its mechanisms of action, lavender may offer a valuable complementary approach in the management of various nervous system disorders, providing an accessible and well-tolerated alternative to conventional treatments [10]. This review examines the clinical evidence supporting lavender's efficacy in neurological applications, emphasizing its potential mechanisms and therapeutic benefits.

2. Human Studies on Lavender and the Nervous System

2.1 Anxiety and Depression

Several clinical trials have assessed the anxiolytic and antidepressant effects of lavender oil. A randomized controlled trial demonstrated that oral administration of lavender oil (Silexan, 80 mg/day) significantly reduced symptoms of generalized anxiety disorder, with efficacy comparable to lorazepam. Additionally, patients reported an improvement in overall well-being without experiencing the common side effects associated with conventional anxiolytic medications, such as sedation and dependency [11-12].

Lavender oil has also been shown to enhance sleep quality, alleviate restlessness, and reduce somatic complaints commonly associated with anxiety disorders. These findings have been reinforced by research on intensive care unit (ICU) patients, where lavender aromatherapy significantly reduced anxiety levels and improved mood. Hemodialysis patients, who frequently experience high levels of stress and psychological distress, also benefited from lavender inhalation therapy, reporting decreased anxiety and improved emotional balance [13].

Moreover, postpartum women suffering from anxiety and depression showed remarkable improvements after undergoing lavender aromatherapy. In a clinical study, the administration of lavender essential oil significantly reduced scores on the Edinburgh Postnatal Depression Scale and the Generalized Anxiety Disorder Scale over a four-week period. These findings suggest that lavender may serve as an effective and natural intervention for postpartum mood disorders, offering an alternative to pharmacological treatments that may not be suitable for breastfeeding mothers [14].

Further studies have explored the neurobiological mechanisms underlying lavender's anxiolytic and antidepressant effects. Research indicates that lavender oil interacts with the gamma-aminobutyric acid (GABA) system, enhancing inhibitory neurotransmission and reducing neuronal excitability. This mechanism is similar to that of benzodiazepines but without the associated side effects, making lavender a promising candidate for long-term anxiety management. Additionally, neuroimaging studies have shown that lavender aromatherapy influences brain regions involved in emotional regulation, such as the amygdala and prefrontal cortex, further supporting its role in alleviating anxiety and depressive symptoms [15]. Despite these promising findings, larger and more rigorous clinical trials are needed to establish standardized dosing guidelines and determine the long-term safety and efficacy of lavender oil for anxiety and depression treatment. Nonetheless, current evidence suggests that lavender holds significant potential as a complementary or standalone therapy for individuals experiencing mood disorders, particularly those seeking natural alternatives to traditional pharmacological treatments.

2.2 Neuroimaging and EEG Studies

Neuroimaging studies have explored the effects of lavender on brain activity, shedding light on its potential mechanisms of action in modulating cognitive and emotional regulation. Functional MRI (fMRI) studies have demonstrated that lavender exposure increases activation in key brain regions, including the orbitofrontal cortex, hippocampus, thalamus, and cerebellum. These areas are associated with memory processing, stress

modulation, and mood regulation, suggesting that lavender may exert its anxiolytic and antidepressant effects through direct neural modulation. Moreover, studies using positron emission tomography (PET) scans have shown changes in metabolic activity in the amygdala, a region heavily involved in fear and emotional responses. The reduction in amygdala activity following lavender inhalation aligns with its anxiolytic properties, indicating that it may help mitigate hyperactivity in stress-related neural circuits. Additionally, changes in cerebral blood flow patterns in response to lavender suggest enhanced connectivity between emotion-regulating centers of the brain [16].

Electroencephalography (EEG) studies have further supported these findings, revealing increased alpha and theta wave activity after lavender exposure. Elevated alpha wave activity is commonly associated with states of wakeful relaxation, reduced anxiety, and improved emotional stability, while theta waves are linked to deep relaxation and enhanced memory consolidation. These EEG patterns align closely with the physiological effects observed in clinical settings, where patients report feelings of calmness and improved sleep following lavender aromatherapy or oral supplementation [17, 18].

Beyond anxiety reduction, EEG data also suggests that lavender may have cognitive-enhancing properties. Some studies indicate that exposure to lavender can modulate brain wave activity in ways that promote attentional control and focus, although findings remain mixed. While some reports highlight improved cognitive performance, others suggest mild impairments in working memory, possibly due to its sedative-like effects [19]. In conclusion, neuroimaging and EEG studies provide compelling evidence that lavender influences brain function in ways that support its therapeutic applications. Future research should focus on expanding our understanding of the long-term neural effects of lavender and optimizing its use in clinical and therapeutic settings.

2.3 Sleep Disorders

Lavender has been widely studied as a natural treatment for insomnia and other sleep disorders, with growing evidence supporting its efficacy in improving sleep quality and reducing sleep disturbances. Randomized controlled trials have demonstrated that lavender oil enhances sleep parameters in various populations, including students, cardiac patients, and menopausal women. These studies report improvements in sleep latency, duration, and overall sleep satisfaction, highlighting lavender's potential as a viable alternative to pharmaceutical sleep aids [20].

In addition to its effectiveness in the general population, lavender has shown promising results in clinical settings involving patients with chronic conditions that impact sleep. A study conducted on benzodiazepine-dependent geriatric patients found that substituting lavender aromatherapy for sedatives helped maintain sleep duration while reducing the need for medication. This suggests that lavender may aid in weaning patients off pharmacological treatments, thereby minimizing the risk of dependency and associated side effects [21].

Mechanistically, lavender's impact on sleep is attributed to its interaction with the central nervous system. Research indicates that lavender enhances gamma-aminobutyric acid (GABA) activity, promoting relaxation and reducing neural excitability. This aligns with EEG findings that show increased alpha and theta wave activity, which are associated with a state of calmness and improved sleep quality. Additionally, lavender has been found to lower cortisol levels, further supporting its role in stress reduction and sleep regulation. The mode of lavender administration also plays a crucial role in its effectiveness. Inhalation of lavender essential oil, through diffusion or pillow sprays, has been widely studied and shown to improve subjective sleep quality. Oral supplementation with lavender oil capsules has similarly yielded positive results, with one study reporting significant improvements in sleep efficiency and reduced nighttime awakenings in patients with mild to moderate anxiety [22].

Despite these promising findings, further research is needed to standardize optimal dosing and administration methods. Larger-scale clinical trials should explore long-term effects, particularly in populations with severe sleep disorders such as insomnia and sleep apnea [23]. Nevertheless, current evidence strongly supports the potential of lavender as a non-pharmacological intervention for sleep improvement, offering a safe and natural alternative for individuals struggling with sleep disturbances.

2.4 Pain Management

Lavender oil has demonstrated significant analgesic effects in clinical settings, offering a promising natural alternative for pain relief. Studies have shown that lavender aromatherapy effectively reduces postoperative pain, lowers the need for opioid analgesics, and alleviates pain associated with various conditions, including migraine, dysmenorrhea, and aphthous ulcers. Its pain-relieving properties are thought to be mediated through its ability to modulate pain perception, reduce inflammation, and induce relaxation by interacting with the central nervous system [24].

Postoperative pain management has been a major area of focus in lavender research. Clinical trials have demonstrated that patients undergoing surgical procedures, including cesarean section and episiotomy recovery, experienced significant pain relief with lavender oil application. Inhalation of lavender aromatherapy post-surgery was associated with reduced pain scores and lower opioid consumption, indicating its potential as an adjunct to conventional analgesics. Additionally, some studies have suggested that lavender may contribute to faster wound healing, further benefiting post-surgical recovery [25].

Beyond postoperative pain, lavender has also shown efficacy in managing chronic pain conditions. Patients with nonspecific neck and lower back pain reported reduced pain intensity and improved mobility following lavender massage therapy. The application of lavender oil in massage has been found to enhance circulation, decrease muscle tension, and promote overall relaxation, making it an effective approach for individuals suffering from musculoskeletal pain [26].

Migraine and headache relief have also been well-documented among lavender's therapeutic benefits. Clinical trials indicate that inhaling lavender oil during the early stages of a migraine attack can significantly reduce the severity and duration of headaches. The mechanism is believed to involve lavender's ability to reduce vascular inflammation, modulate serotonin levels, and enhance parasympathetic nervous system activity, contributing to its pain-relieving effects [27]. Additionally, lavender has been explored as a treatment for menstrual pain. Studies on women with primary dysmenorrhea have reported that lavender aromatherapy and abdominal massage with lavender oil effectively reduce pain intensity and improve overall well-being. Its muscle relaxant properties may play a key role in alleviating uterine contractions and discomfort associated with menstrual pain [28].

Overall, the evidence supports lavender as a viable complementary therapy for pain management. While further research is needed to establish standardized dosages and administration methods, its effectiveness in reducing pain and improving patient comfort highlights its potential as a natural, non-invasive pain relief option in both acute and chronic conditions.

2.5 Cognitive Function and Memory

The effects of lavender on cognitive performance have yielded mixed results, with research indicating both potential benefits and limitations depending on the specific cognitive domain being assessed. Some studies have reported mood enhancement and improved concentration, leading to reduced errors in mathematical and logical reasoning tasks after lavender exposure. The calming effects of lavender are believed to contribute to better focus and stress reduction, which can enhance cognitive performance in certain contexts. However, other studies have suggested that lavender may have sedative-like

properties that could impair working memory and reaction times. For instance, in tasks requiring quick decision-making and sustained attention, participants exposed to lavender aromatherapy showed slower response times, indicating that while lavender promotes relaxation, it may also dampen cognitive alertness in some individuals. This raises questions about its optimal use, particularly in activities requiring high levels of cognitive engagement [29].

Research on elderly populations, particularly those with neurodegenerative conditions such as dementia, has yielded promising results. A study on dementia patients found that aromatherapy with lavender and other essential oils improved cognitive orientation, emotional state, and overall well-being. Lavender's ability to modulate the limbic system may contribute to reducing agitation and improving mood stability in patients with Alzheimer's disease and related disorders. Some studies suggest that consistent exposure to lavender may enhance sleep quality in patients with dementia, which in turn positively impacts cognitive function [30-31].

Beyond neurodegenerative conditions, preliminary research has explored lavender's effects on learning and memory consolidation. Some findings suggest that lavender may influence hippocampal activity, a brain region crucial for memory processing. In animal models, lavender extract has demonstrated neuroprotective effects, but more research is needed to determine whether similar benefits extend to human cognition. Additionally, EEG studies have shown increased alpha and theta wave activity after lavender exposure, which is associated with relaxation and memory consolidation [32].

Despite the mixed findings, lavender's potential as a complementary therapy for cognitive enhancement remains an area of interest. Further studies are needed to clarify its role in cognitive function across different populations, refine its application methods, and establish the appropriate balance between its calming effects and cognitive performance outcomes. Future research should focus on standardizing dosages and evaluating long-term effects to better understand how lavender can be integrated into cognitive health interventions.

3. Mechanisms of Action

The neurological effects of lavender are believed to be mediated through multiple mechanisms, which contribute to its anxiolytic, antidepressant, neuroprotective, and cognitive-modulating properties. These mechanisms are supported by biochemical, neuroimaging, and behavioral studies. One of the primary ways lavender exerts its effects is through the modulation of the limbic system, particularly the amygdala and hippocampus. These brain regions are critical for processing emotions, stress responses, and memory regulation. Functional imaging studies have demonstrated that lavender exposure influences activity in these areas, reducing hyperactivity in stress-related circuits and promoting a sense of calmness and emotional stability. This modulation may explain its efficacy in reducing symptoms of anxiety and depression, as well as its potential benefits for cognitive function in neurodegenerative conditions [15].

Lavender also interacts with gamma-aminobutyric acid (GABA) receptors, which are responsible for inhibitory neurotransmission in the central nervous system. By enhancing GABAergic activity, lavender promotes relaxation, reduces neuronal excitability, and mitigates excessive nervous system arousal. This mechanism is like that of benzodiazepines, though without the risk of dependence and sedation commonly associated with pharmaceutical anxiolytics [33]. EEG studies have shown increased alpha and theta wave activity following lavender exposure, reinforcing its sedative and anxiolytic effects. Additionally, lavender has been shown to reduce cortisol levels, the primary stress hormone involved in the body's response to psychological and physiological stressors [34]. Chronic stress and elevated cortisol levels are linked to anxiety, depression, and cognitive impairment. Studies indicate that inhalation or oral administration of lavender essential oil can lead to significant reductions in cortisol, further supporting its role as a stress modulator and mood stabilizer [35].

Beyond its impact on neurotransmission and stress hormones, lavender possesses notable antioxidant and anti-inflammatory properties. These properties contribute to neuroprotection by reducing oxidative stress, which is implicated in neurodegenerative diseases such as Alzheimer's and Parkinson's [36]. Lavender's bioactive compounds, including linalool and linalyl acetate, have been shown to counteract neuroinflammation and enhance neuronal resilience against damage caused by free radicals and toxic insults [37]. Furthermore, emerging research suggests that lavender may influence dopaminergic and serotonergic neurotransmission, pathways that play key roles in mood regulation and cognitive function. By modulating these neurotransmitter systems, lavender may have antidepressant effects and potential cognitive-enhancing properties, though further research is needed to clarify these interactions [38].

Overall, the multifaceted mechanisms of action underlying lavender's neurological effects provide a strong basis for its therapeutic applications. Future studies should aim to further elucidate these mechanisms, optimize dosing strategies, and explore its potential in combination with conventional treatments for neurological and psychiatric disorders.

4. Safety and Adverse Effects

Lavender oil is generally considered safe for therapeutic use, but some adverse effects have been reported, particularly with prolonged or excessive exposure. While many individuals tolerate lavender well, certain populations may be more susceptible to its potential side effects. One of the most commonly reported adverse effects is mild gastrointestinal discomfort, including nausea, bloating, and dyspepsia, particularly when lavender oil is consumed orally. These symptoms are generally self-limiting and resolve with dosage adjustments. Inhalation or topical use is less likely to cause gastrointestinal issues, making these administration routes preferable for individuals prone to digestive sensitivity [39].

Allergic reactions to lavender oil have also been documented, especially in individuals with sensitivities to essential oils. Contact dermatitis, characterized by skin redness, itching, and irritation, can occur when lavender oil is applied topically, particularly in undiluted form. Those with known allergies to Lamiaceae family plants should exercise caution when using lavender-based products. Conducting a patch test before topical application can help minimize the risk of allergic reactions [40].

Concerns regarding potential hormonal effects have been raised, particularly in relation to prepubertal gynecomastia in boys. Some case studies suggest that repeated topical application of lavender and tea tree oil may have estrogenic and antiandrogenic properties, potentially leading to hormonal imbalances. However, more extensive clinical studies are needed to confirm these effects and determine their clinical significance. Until further evidence is available, it may be advisable to avoid frequent topical use of lavender oil in children and adolescents [41].

Another area of caution is the use of lavender oil during pregnancy and breastfeeding. While no severe teratogenic effects have been reported, lavender's potential influence on hormonal pathways raises concerns about its safety in pregnant individuals. Some experts recommend avoiding oral consumption of lavender oil during pregnancy due to its potential emmenagogue (menstruation-stimulating) properties. Similarly, breastfeeding mothers should use lavender with caution, as the effects on infant development remain insufficiently studied [42].

Despite these concerns, lavender oil has no known potential for abuse or dependency, making it a safer alternative to many pharmacological treatments for anxiety, sleep disorders, and pain. However, it should always be used in moderation and under the guidance of a healthcare professional, particularly in vulnerable populations such as pregnant women, children, and individuals with pre-existing medical conditions [43]. Finally, while lavender is generally well-tolerated and safe for most users, awareness of potential

side effects and contraindications is essential for its responsible use. Future research should focus on clarifying the long-term safety profile of lavender, particularly in populations with heightened sensitivity to hormonal influences.

5. Conclusion

Clinical evidence supports the use of lavender essential oil for anxiety, sleep disorders, pain management, and mood regulation. However, further high-quality studies with larger sample sizes and standardized methodologies are needed to confirm its long-term efficacy and safety. Future research should focus on elucidating the precise mechanisms of action and optimizing therapeutic formulations for clinical applications.

Funding: None.

Research Ethics Committee Approval: None.

Acknowledgments: None.

Conflicts of Interest: None.

Supplementary Materials: None.

References

1. Batiha GE, Teibo JO, Wasef L, Shaheen HM, Akomolafe AP, Teibo TKA, Al-Kuraishy HM, Al-Garbeeb AI, Alexiou A, Papadakis M. A review of the bioactive components and pharmacological properties of *Lavandula* species. *Naunyn Schmiedebergs Arch Pharmacol*. 2023 May;396(5):877-900. doi: 10.1007/s00210-023-02392-x. Epub 2023 Feb 11. PMID: 36773055; PMCID: PMC10079719.
2. Koulivand PH, Khaleghi Ghadiri M, Gorji A. Lavender and the nervous system. *Evid Based Complement Alternat Med*. 2013;2013:681304. doi: 10.1155/2013/681304. Epub 2013 Mar 14. PMID: 23573142; PMCID: PMC3612440.
3. Bavarsad NH, Bagheri S, Kourosh-Arami M, Komaki A. Aromatherapy for the brain: Lavender's healing effect on epilepsy, depression, anxiety, migraine, and Alzheimer's disease: A review article. *Heliyon*. 2023 Jul 20;9(8):e18492. doi: 10.1016/j.heliyon.2023.e18492. PMID: 37554839; PMCID: PMC10404968.
4. Donelli D, Antonelli M, Bellinazzi C, Gensini GF, Firenzuoli F. Effects of lavender on anxiety: A systematic review and meta-analysis. *Phytomedicine*. 2019 Dec;65:153099. doi: 10.1016/j.phymed.2019.153099. Epub 2019 Sep 26. PMID: 31655395.
5. Fung TKH, Lau BWM, Ngai SPC, Tsang HWH. Therapeutic Effect and Mechanisms of Essential Oils in Mood Disorders: Interaction between the Nervous and Respiratory Systems. *Int J Mol Sci*. 2021 May 3;22(9):4844. doi: 10.3390/ijms22094844. PMID: 34063646; PMCID: PMC8125361.
6. Cui J, Li M, Wei Y, Li H, He X, Yang Q, Li Z, Duan J, Wu Z, Chen Q, Chen B, Li G, Ming X, Xiong L, Qin D. Inhalation Aromatherapy via Brain-Targeted Nasal Delivery: Natural Volatiles or Essential Oils on Mood Disorders. *Front Pharmacol*. 2022 Apr 12;13:860043. doi: 10.3389/fphar.2022.860043. PMID: 35496310; PMCID: PMC9041268.
7. Firoozeei TS, Feizi A, Rezaeizadeh H, Zargarani A, Roohafza HR, Karimi M. The antidepressant effects of lavender (*Lavandula angustifolia* Mill.): A systematic review and meta-analysis of randomized controlled clinical trials. *Complement Ther Med*. 2021 Jun;59:102679. doi: 10.1016/j.ctim.2021.102679. Epub 2021 Feb 4. PMID: 33549687.
8. Lin PW, Chan WC, Ng BF, Lam LC. Efficacy of aromatherapy (*Lavandula angustifolia*) as an intervention for agitated behaviours in Chinese older persons with dementia: a cross-over randomized trial. *Int J Geriatr Psychiatry*. 2007 May;22(5):405-10. doi: 10.1002/gps.1688. PMID: 17342790.
9. Amitani H, Nishi R, Sagiya K, Fukumoto T, Funakoshi K, Takayanagi N, Watanabe H, Hirose M, Tagawa K, Ota K, Ito YM, Asakawa A. The effect of lavender aroma for anxiety disorder: a study protocol for a multicenter, double-masked, randomized, placebo-controlled clinical trial. *BMC Complement Med Ther*. 2023 Nov 6;23(1):397. doi: 10.1186/s12906-023-04231-1. PMID: 37932761; PMCID: PMC10626714.
10. López V, Nielsen B, Solas M, Ramírez MJ, Jäger AK. Exploring Pharmacological Mechanisms of Lavender (*Lavandula angustifolia*) Essential Oil on Central Nervous System Targets. *Front Pharmacol*. 2017 May 19;8:280. doi: 10.3389/fphar.2017.00280. PMID: 28579958; PMCID: PMC5437114.
11. Dold M, Bartova L, Volz HP, Seifritz E, Möller HJ, Schläpke S, Kasper S. Efficacy of Silexan in patients with anxiety disorders: a meta-analysis of randomized, placebo-controlled trials. *Eur Arch Psychiatry Clin Neurosci*. 2023 Oct;273(7):1615-1628. doi: 10.1007/s00406-022-01547-w. Epub 2023 Jan 30. PMID: 36717399; PMCID: PMC10465640.
12. Kasper S, Gastpar M, Müller WE, Volz HP, Möller HJ, Schläpke S, Dienel A. Lavender oil preparation Silexan is effective in generalized anxiety disorder—a randomized, double-blind comparison to placebo and paroxetine. *Int J Neuropsychopharmacol*. 2014 Jun;17(6):859-69. doi: 10.1017/S1461145714000017. Epub 2014 Jan 23. PMID: 24456909.

13. Karadag E, Samancioglu S, Ozden D, Bakir E. Effects of aromatherapy on sleep quality and anxiety of patients. *Nurs Crit Care*. 2017 Mar;22(2):105-112. doi: 10.1111/nicc.12198. Epub 2015 Jul 27. PMID: 26211735.
14. Kianpour M, Mansouri A, Mehrabi T, Asghari G. Effect of lavender scent inhalation on prevention of stress, anxiety and depression in the postpartum period. *Iran J Nurs Midwifery Res*. 2016 Mar-Apr;21(2):197-201. doi: 10.4103/1735-9066.178248. PMID: 27095995; PMCID: PMC4815377.
15. López V, Nielsen B, Solas M, Ramírez MJ, Jäger AK. Exploring Pharmacological Mechanisms of Lavender (*Lavandula angustifolia*) Essential Oil on Central Nervous System Targets. *Front Pharmacol*. 2017 May 19;8:280. doi: 10.3389/fphar.2017.00280. PMID: 28579958; PMCID: PMC5437114.
16. Sayorwan W, Siripornpanich V, Piriapunyaporn T, Hongratanaworakit T, Kotchabhakdi N, Ruangrunsi N. The effects of lavender oil inhalation on emotional states, autonomic nervous system, and brain electrical activity. *J Med Assoc Thai*. 2012 Apr;95(4):598-606. PMID: 22612017.
17. Lagopoulos J, Xu J, Rasmussen I, Vik A, Malhi GS, Eliassen CF, Arntsen IE, Saether JG, Hollup S, Holen A, Davanger S, Ellingsen Ø. Increased theta and alpha EEG activity during nondirective meditation. *J Altern Complement Med*. 2009 Nov;15(11):1187-92. doi: 10.1089/acm.2009.0113. PMID: 19922249.
18. Klimesch W. EEG alpha and theta oscillations reflect cognitive and memory performance: a review and analysis. *Brain Res Brain Res Rev*. 1999 Apr;29(2-3):169-95. doi: 10.1016/s0165-0173(98)00056-3. PMID: 10209231.
19. Afghan R, Heysieattalab S, Zangbar HS, Ebrahimi-Kalan A, Jafari-Koshki T, Samadzadehaghdam N. Lavender Essential Oil Inhalation Improves Attentional Shifting and Accuracy: Evidence from Dynamic Changes of Cognitive Flexibility and Power Spectral Density of Electroencephalogram Signals. *J Med Signals Sens*. 2024 Apr 18;14:12. doi: 10.4103/jmss.jmss_57_23. PMID: 38993201; PMCID: PMC11111129.
20. Chen TY, Hiyama A, Muramatsu M, Hinotsu A. The Effect of Lavender on Sleep Quality in Individuals Without Insomnia: A Systematic Review. *Holist Nurs Pract*. 2022 Jul-Aug 01;36(4):223-231. doi: 10.1097/HNP.0000000000000528. PMID: 35708558.
21. Luo J, Jiang W. A critical review on clinical evidence of the efficacy of lavender in sleep disorders. *Phytother Res*. 2022 Jun;36(6):2342-2351. doi: 10.1002/ptr.7448. Epub 2022 Apr 12. PMID: 35412693.
22. Ren YL, Chu WW, Yang XW, Xin L, Gao JX, Yan GZ, Wang C, Chen YN, Xie JF, Spruyt K, Lin JS, Hou YP, Shao YF. Lavender improves sleep through olfactory perception and GABAergic neurons of the central amygdala. *J Ethnopharmacol*. 2025 Jan 30;337(Pt 3):118942. doi: 10.1016/j.jep.2024.118942. Epub 2024 Oct 18. PMID: 39426576.
23. Chang JL, Goldberg AN, Alt JA, Mohammed A, Ashbrook L, Auckley D, Ayappa I, Bakhtiar H, Barrera JE, Bartley BL, Billings ME, Boon MS, Bosschieter P, Braverman I, Brodie K, Cabrera-Muffly C, Caesar R, Cahali MB, Cai Y, Cao M, Capasso R, Caples SM, Chahine LM, Chang CP, Chang KW, Chaudhary N, Cheong CSJ, Chowdhuri S, Cistulli PA, Claman D, Collen J, Coughlin KC, Creamer J, Davis EM, Dupuy-McCauley KL, Durr ML, Dutt M, Ali ME, Elkassabany NM, Epstein LJ, Fiala JA, Freedman N, Gill K, Gillespie MB, Golisch L, Gooneratne N, Gottlieb DJ, Green KK, Gulati A, Gurubhagavatula I, Hayward N, Hoff PT, Hoffmann OMG, Holfinger SJ, Hsia J, Huntley C, Huoh KC, Huyett P, Inala S, Ishman SL, Jella TK, Jobanputra AM, Johnson AP, Junna MR, Kado JT, Kaffenberger TM, Kapur VK, Kezirian EJ, Khan M, Kirsch DB, Kominsky A, Kryger M, Krystal AD, Kushida CA, Kuzniar TJ, Lam DJ, Lettieri CJ, Lim DK, Lin HC, Liu SYC, MacKay SG, Magalang UJ, Malhotra A, Mansukhani MP, Maurer JT, May AM, Mitchell RB, Mokhlesi B, Mullins AE, Nada EM, Naik S, Nokes B, Olson MD, Pack AI, Pang EB, Pang KP, Patil SP, Van de Perck E, Piccirillo JF, Pien GW, Piper AJ, Plawecki A, Quigg M, Ravesloot MJL, Redline S, Rothenberg BW, Ryden A, Sarmiento KF, Sbeih F, Schell AE, Schmickl CN, Schotland HM, Schwab RJ, Seo J, Shah N, Shelgikar AV, Shochat I, Soose RJ, Steele TO, Stephens E, Stepnowsky C, Strohl KP, Sutherland K, Suurna MV, Thaler E, Thapa S, Vanderveken OM, de Vries N, Weaver EM, Weir ID, Wolfe LF, Woodson BT, Won CHJ, Xu J, Yalamanchi P, Yaremchuk K, Yeghiazarians Y, Yu JL, Zeidler M, Rosen IM. International Consensus Statement on Obstructive Sleep Apnea. *Int Forum Allergy Rhinol*. 2023 Jul;13(7):1061-1482. doi: 10.1002/alr.23079. Epub 2023 Mar 30. PMID: 36068685; PMCID: PMC10359192.
24. Grabnar M, Roach MJ, Abd-Elseyed A, Kim C. Impact of Lavender on Pain and Anxiety Levels Associated With Spine Procedures. *Ochsner J*. 2021 Winter;21(4):358-363. doi: 10.31486/toj.21.0013. PMID: 34984050; PMCID: PMC8675619.
25. Abedian S, Abedi P, Jahanfar S, Irvani M, Zahedian M. The effect of Lavender on pain and healing of episiotomy: A systematic review. *Complement Ther Med*. 2020 Sep;53:102510. doi: 10.1016/j.ctim.2020.102510. Epub 2020 Jul 28. PMID: 33066851.
26. Mizrak Sahin B, Culha I, Gursoy E, Yalcin OT. Effect of Massage With Lavender Oil on Postoperative Pain Level of Patients Who Underwent Gynecologic Surgery: A Randomized, Placebo-Controlled Study. *Holist Nurs Pract*. 2021 Jul-Aug 01;35(4):221-229. doi: 10.1097/HNP.0000000000000400. PMID: 32657903.
27. Sasannejad P, Saeedi M, Shoeibi A, Gorji A, Abbasi M, Foroughipour M. Lavender essential oil in the treatment of migraine headache: a placebo-controlled clinical trial. *Eur Neurol*. 2012;67(5):288-91. doi: 10.1159/000335249. Epub 2012 Apr 17. PMID: 22517298.
28. Nikjou R, Kazemzadeh R, Rostamnegad M, Moshfegi S, Karimollahi M, Salehi H. The Effect of Lavender Aromatherapy on the Pain Severity of Primary Dysmenorrhea: A Triple-blind Randomized Clinical Trial. *Ann Med Health Sci Res*. 2016 Jul-Aug;6(4):211-215. doi: 10.4103/amhsr.amhsr_527_14. PMID: 28480095; PMCID: PMC5405632.
29. Malloggi E, Menicucci D, Cesari V, Frumento S, Gemignani A, Bertoli A. Lavender aromatherapy: A systematic review from essential oil quality and administration methods to cognitive enhancing effects. *Appl Psychol Health Well Being*. 2022 May;14(2):663-690. doi: 10.1111/aphw.12310. Epub 2021 Oct 5. PMID: 34611999; PMCID: PMC9291879.
30. Ball EL, Owen-Booth B, Gray A, Shenkin SD, Hewitt J, McCleery J. Aromatherapy for dementia. *Cochrane Database Syst Rev*. 2020 Aug 19;8(8):CD003150. doi: 10.1002/14651858.CD003150.pub3. PMID: 32813272; PMCID: PMC7437395.

31. Jimbo D, Kimura Y, Taniguchi M, Inoue M, Urakami K. Effect of aromatherapy on patients with Alzheimer's disease. *Psychogeriatrics*. 2009 Dec;9(4):173-9. doi: 10.1111/j.1479-8301.2009.00299.x. PMID: 20377818.
32. Avola R, Furnari AG, Graziano ACE, Russo A, Cardile V. Management of the Brain: Essential Oils as Promising Neuroinflammation Modulator in Neurodegenerative Diseases. *Antioxidants (Basel)*. 2024 Jan 31;13(2):178. doi: 10.3390/antiox13020178. PMID: 38397776; PMCID: PMC10886016.
33. Wang ZJ, Heinbockel T. Essential Oils and Their Constituents Targeting the GABAergic System and Sodium Channels as Treatment of Neurological Diseases. *Molecules*. 2018 May 2;23(5):1061. doi: 10.3390/molecules23051061. PMID: 29724056; PMCID: PMC6099651.
34. Baumgardner DJ. Lavender Aromatherapy: Ready for Prime-Time Office Use? *J Patient Cent Res Rev*. 2020 Jan 27;7(1):5-7. PMID: 32002442; PMCID: PMC6988708.
35. Yoo O, Park SA. Anxiety-Reducing Effects of Lavender Essential Oil Inhalation: A Systematic Review. *Healthcare (Basel)*. 2023 Nov 17;11(22):2978. doi: 10.3390/healthcare11222978. PMID: 37998470; PMCID: PMC10671255.
36. Faridzadeh A, Salimi Y, Ghasemirad H, Kargar M, Rashtchian A, Mahmoudvand G, Karimi MA, Zerangian N, Jahani N, Masoudi A, Sadeghian Dastjerdi B, Salavatizadeh M, Sadeghsalehi H, Deravi N. Neuroprotective Potential of Aromatic Herbs: Rosemary, Sage, and Lavender. *Front Neurosci*. 2022 Jun 28;16:909833. doi: 10.3389/fnins.2022.909833. PMID: 35873824; PMCID: PMC9297920.
37. Shin YK, Seol GH. Effects of linalyl acetate on oxidative stress, inflammation and endothelial dysfunction: can linalyl acetate prevent mild cognitive impairment? *Front Pharmacol*. 2023 Jul 28;14:1233977. doi: 10.3389/fphar.2023.1233977. PMID: 37576815; PMCID: PMC10416234.
38. Dos Santos ÉRQ, Maia JGS, Fontes-Júnior EA, do Socorro Ferraz Maia C. Linalool as a Therapeutic and Medicinal Tool in Depression Treatment: A Review. *Curr Neuropharmacol*. 2022;20(6):1073-1092. doi: 10.2174/1570159X19666210920094504. PMID: 34544345; PMCID: PMC9886818.
39. Kajjari S, Joshi RS, Hugar SM, Gokhale N, Meharwade P, Uppin C. The Effects of Lavender Essential Oil and its Clinical Implications in Dentistry: A Review. *Int J Clin Pediatr Dent*. 2022 May-Jun;15(3):385-388. doi: 10.5005/jp-journals-10005-2378. PMID: 35991803; PMCID: PMC9357533.
40. Bingham LJ, Tam MM, Palmer AM, Cahill JL, Nixon RL. Contact allergy and allergic contact dermatitis caused by lavender: A retrospective study from an Australian clinic. *Contact Dermatitis*. 2019 Jul;81(1):37-42. doi: 10.1111/cod.13247. Epub 2019 Apr 16. PMID: 30779160.
41. Henley DV, Lipson N, Korach KS, Bloch CA. Prepubertal gynecomastia linked to lavender and tea tree oils. *N Engl J Med*. 2007 Feb 1;356(5):479-85. doi: 10.1056/NEJMoa064725. PMID: 17267908.
42. Vidal-García E, Vallhonrat-Bueno M, Pla-Consuegra F, Orta-Ramírez A. Efficacy of Lavender Essential Oil in Reducing Stress, Insomnia, and Anxiety in Pregnant Women: A Systematic Review. *Healthcare (Basel)*. 2024 Dec 5;12(23):2456. doi: 10.3390/healthcare12232456. PMID: 39685078; PMCID: PMC11641599.
43. Despite these concerns, lavender oil has no known potential for abuse or dependency, making it a safer alternative to many pharmacological treatments for anxiety, sleep disorders, and pain. However, it should always be used in moderation and under the guidance of a healthcare professional, particularly in vulnerable populations such as pregnant women, children, and individuals with pre-existing medical conditions.